

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Characterizing Chemical Exposure Trends from NHANES Urinary Biomonitoring Data

Zachary Stanfield, R. Woodrow Setzer, Victoria Hull, Risa R. Sayre, Kristin K. Isaacs, and John F. Wambaugh

Table of Contents

Figure S1. Heatmap of the 2-year NHANES cohorts (columns) in which each of the 151 metabolites (rows; chemical names not shown) were measured in NHANES. The metabolites are sorted from top to bottom by oldest cohort with measurement data then by most cohorts with data. A table representation of this figure, with metabolites in the same order, can be found in Excel Table S14.

Figure S2. Distributions of the number of survey participants and measurements below the limit of detection (LOD). Each point represents a metabolite-cohort pair (single metabolite measured in a single NHANES cohort), depicting the fraction of observations below the LOD on the y-axis and the number of individual survey participants on the x-axis. The histograms on the top and to the right of the scatterplot represent the count distribution for all metabolite-cohort pairs of the number of survey participants and the fraction of urine measurements with a concentration below the LOD, respectively. No chemical filtering was applied prior to making this plot, meaning all 151 metabolites have a point in this figure for each cohort in which they were measured in urine by NHANES. The data used to generate this plot can be found in Excel Table S15.

Additional File- Excel Document

Supplemental Figures

Metabolite Data Availability Across NHANES Cohorts




Figure S1. Heatmap of the 2-year NHANES cohorts (columns) in which each of the 151 metabolites (rows; chemical names not shown) were measured in NHANES. The metabolites are sorted from top to bottom by oldest cohort with measurement data then by most cohorts with data. A table representation of this figure, with metabolites in the same order, can be found in Excel Table S14.




Figure S2. Distributions of the number of survey participants and measurements below the limit of detection (LOD). Each point represents a metabolite-cohort pair (single metabolite measured in a single NHANES cohort), depicting the fraction of observations below the LOD on the y-axis and the number of individual survey participants on the x-axis. The histograms on the top and to the right of the scatterplot represent the count distribution for all metabolite-cohort pairs of the number of survey participants and the fraction of urine measurements with a concentration below the LOD, respectively. No chemical filtering was applied prior to making this plot, meaning all 151 metabolites have a point in this figure for each cohort in which they were measured in urine by NHANES. The data used to generate this plot can be found in Excel Table S15.